



# Let's Talk TB:

A Supplement to GP CLINICS

## **Chapter 12:**

What counseling and support do patients with Tuberculosis need?

Authors: Indira Behara, MBBS, MPH; Karishma Saran, MSc; Aasit G. Nanavati, MPH; Deepti Chavan Musale, Patient Advocate



- Have you ever walked into your GP's clinic, waited in line for 30 minutes, spent about 5 minutes with your doctor and left with a prescription in hand?
- Unfortunately, due to the extremely busy schedules of our GPs, the average consultation time lasts barely a few minutes
  - During this time, very few questions are asked of the patient



- A crucial step missing in this process is that of 'patient counseling'
  - In India, the patient-provider relationship is often one where there is unparalleled respect for the physician
  - The patient often has blind faith in their diagnosis and subsequent treatment guidelines
- Due to this asymmetric relationship and rushed consultation, patients do not always receive the adequate education and counseling they deserve



- Patient education is defined as follows:
  - Education to help patients and their families understand the disease and treatment, cooperate with healthcare providers, live healthily, and maintain or improve their quality of life



- In the case of a disease such as tuberculosis (TB), patient education along with appropriate counseling (i.e. enabling them to cope with the stress and take personal decisions related to the disease) is the need of the hour
- TB in India is plagued by a number of challenges
  - Some of these are related to the patient's inaccurate or incomplete knowledge, incorrect behavior beliefs and stigma-generating attitudes
  - Patient counseling can therefore play a crucial to counter these problems



- The Revised National Tuberculosis Programme (RNTCP)
   of the Ministry of Health and Family Welfare (MoHFW)
   also recognizes the urgent need to scale up counseling
   and psychosocial support services for patients
- India's ambitious National Strategic Plan (NSP) for TB prevention and control (2012 – 2017) highlights that patient counseling through a professional counselor is a missing link in the programme
- It attributes counseling to be a critical intervention that can enhance treatment adherence and improve treatment outcomes



- The journey of a TB patient from the time of the onset of symptoms to treatment completion is not an easy one
- Many of the factors that increase a person's vulnerability to TB or reduce their access to services to prevent, diagnose and treat TB are associated with their ability to realize their human rights



- Deepti Chavan Musale (a former TB patient) highlights that:
  - 'a TB patient has to deal with a lot of things including the side effects of the medicines, stigma, discrimination etc. Additionally the duration of treatment is very long and fighting TB is not only a physical battle but one that requires you to be very strong mentally too.'



- In the case of TB, adherence to treatment is vital
- Inability to comply can lead to advanced stages of TB, such as multi drug resistant MDR TB
- TB drugs come with a host of side-effects and counseling helps patients become aware of the effects that they might face and the importance of compliance



- Some of the side effects to commonly used TB drugs may include
  - Feeling sick or dizzy, skin rashes, pins and needles and flu like symptoms
  - Become worse (e.g. jaundice) in the case of second line TB drugs used to cure MDR TB



- Along with the potential of infecting families and communities, there is a deep-rooted stigma attached to a TB patient, which often prevents them from accessing services
- Any gap in timely and accurate diagnosis and treatment perpetuates TB transmission, despite the widespread availability of effective and inexpensive treatment



- The principal effects of stigma in developing countries are social isolation of patients
  - Outside the family (where the person may be avoided by former friends and acquaintances)
  - Inside the family where the patient may be forced to eat and sleep separately



- Patient and family counseling plays an important role in addressing the above issues
  - Also plays a role in helping patients deal with the financial burden and nutritional barriers to cure TB successfully



#### Patient Counseling and Support:

#### when and how

- Psychological distress such as depression and other mental co-morbidities have often been associated with TB
  - With rates of mental illness reaching as high as 70% among TB patients
- Resulting chronic stress can impact a TB patient
  - 1. By impairing his or her immune system
  - 2. Affect his or her ability to adhere to treatment



## Patient Counseling and Support: when and how

 With challenges related to adherence to TB treatment resulting in a number of undesirable consequences, it is important to understand the contributing factors and the role for counseling



## Patient Counseling and Support: when and how

- A critical review by Sabaté describes 5 interacting factors
  - 1. Socio-economic
  - 2. Health system
  - 3. Condition related
  - 4. Therapy related
  - 5. Patient related



#### Counselling

- Can take on the form of simple information sharing or more specialized services directed toward improving outcome expectations for the illness
- Can encourage patient self-efficacy and motivation to achieve a complete cure
- Can be used to address some of these contributing factors



## Patient Counseling and Support: when and how

- There are a number of tool-kits that have been created to train counsellors for patient counselling
- It is important for the physician to also understand and value this component of TB care



## Patient Counseling and Support:

#### when and how

Broadly summarizing the wealth of information cited, it is important to remember:

- The objectives behind counselling
- The type of counselling setting suitable to the patient
- The stage of the patient's care pathway



#### The Objective

- As per the 'Patient's Charter for Tuberculosis
   Care': patients have the right to Care, Dignity,
   Information, Choice and Confidence
- The objective of counselling is to meet the needs and ensure the rights of the patient
- The objective is also to support the patient where possible to overcome barriers to successful treatment



#### The Type

 Counselling can take place through providers adequately trained to identify the needs of the patient, through community, engagements, support groups, and individual counselling



#### The Stage

- Before diagnosis is confirmed, the patient must be informed of and understand all relevant information about
  - The disease
  - Its symptoms
  - The tests
  - The need for follow-up and treatment
  - That it is treatable and curable



#### Diagnosis

- Once diagnosis is confirmed and treatment is to be initiated
  - The patient must understand the regimen and its duration
  - The details of adverse drug reactions and what action to take if they encounter them
  - The importance of adherence and nutrition
  - The importance of infection control



#### Family Support

- It is equally important to enlist family support (if consented to) for the patient to successfully complete treatment
- Once TB is diagnosed, it is important for the patient to also undergo HIV testing and testing for other comorbid conditions such as diabetes
- These tests also require counselling and education



#### **Treatment**

- After treatment is underway (along with monitoring physical and biological parameters in order to ensure adherence, motivation and psychological well-being) it is important that the physician understands the health needs of the patients
  - Including the need to be referred to trained counsellors, community level support groups or mental health services if needed
- It is important therefore to acknowledge the need for a patient centered approach to TB care and understand its benefits



#### Taking the patient centered approach

• The 'Patients Charter for TB Care' states that each patient has a "right to receive timely, concise, and clear description of the medical condition, with diagnosis, prognosis (an opinion as to the likely future course of the illness), and treatment proposed, with communication of common risks and appropriate alternatives



#### Counsellors

- It is this information (which is often overwhelming for the patient) where counsellors can ensure:
  - The patient has a foundation in understanding the illness
  - The critical importance of treatment adherence
- Patient programmes integrating counselling services can also significantly improve treatment adherence and the use of a multimodal approach is more effective in improving adherence than stand-alone programmes



## Implementing a model

- An integral measure in ensuring these patients receive the care they need and continue with their treatment regimen is providing counselling services to strengthen the support system throughout their treatment timeframe
- The model of implementing counselling as a supplement to treatment is not a novel technique
  - Has been successfully implemented for diseases such as HIV/AIDS



#### Notes from a MDR-TB Patient Advocate

- •Patients constantly need someone to motivate them and help them sail through this phase of life
- •A counsellor can guide the patients and advise them on how to deal with their day-to-day problems
- Patients need a good listener who will hear them out and guide them appropriately
- •A patient needs hope, hope that she/he will come out of all this and hope that there will be a better future. Who better than a counsellor can give her/him that?

Watch "Deepti's Story : From Survivor to a Champion" on YouTube - https://youtu.be/hvaormVq0vE



#### **RNTCP**

 Recognizing the importance of providing patient support services that include counselling, the RNTCP has also undertaken a project for providing integrated counselling services to TB patients



#### Project Axshaya

- In May of 2014, in partnership with Population Services International, Project Axshaya was implemented to provide facility and home based counselling for DR-TB patients across 28 districts with the help of 30 DR-TB trained counsellors
- Project Axshaya is implementing strategies that will increase capacity for healthcare providers and ensure patients are equipped with the knowledge to maintain treatment



#### Project Saksham

- Another example of a point-of-care and communitybased counselling is the Saksham project
- Saksham, is implemented by the Tata Institute for Social Sciences (TISS) and was developed to strengthen human and institutional capacities of the national health system in the field of HIV/AIDS counselling
- Counselling services for HIV/AIDS patients have ensured patients understand the disease as well as limit risky behaviors which can increase the spread of the disease



#### Project Saksham

 Owing to the success of Project Saksham, TISS is leading the way in integrating counselling services for TB patients and families, building capacity through training healthcare providers and ensuring patients receive supportive supervised care while being treated for TB



#### Successful models

- It is evident that scaling successful models such as the ones implemented by Project Saksham and Project Axshaya are essential in ensuring TB patients are able and empowered to adhere to treatment guidelines
- Implementing this approach will enable the patient to access the best possible care
- Creating an environment which is supportive of TB patients is integral in addressing the societal problems faced by patients
  - Will strengthen the patients resolve while going through the treatment process



#### Holistic Approach

- TB patients continue to experience difficulties in navigating the path from diagnosis to cure and find it personally distressing
- With the help of a dedicated counselor, the journey to a TB free life would be much easier
  - Is something which is not too difficult to provide
- Rather than focusing on just the disease, it is imperative to treat the patient as a whole



#### Holistic Approach

 We must use patient centered approaches, and recognize that all TB patients deserve a minimum package of holistic TB care services that are not restricted to diagnosis and pharmacological treatment, but include counselling and support services as well



#### Key take away messages for healthcare providers

ALL TB PATIENTS DESERVE A MINIMUM PACKAGE OF COUNSELLING AND TB SERVICES

- Use good communication principles when counselling all patients
- Explain the details of TB:
  - -What TB is?
  - -How the disease spreads?
  - -Signs and symptoms
  - -Consequences of not following treatment guidelines
  - -Why treatment is long and why completion of treatment is critical?
  - -Likely adverse events during therapy
  - -Cost involved in treatment and what free/public services are available to patients
- Identify TB diagnosis and treatment services available to the patient at all government health facilities free of cost
- Explain that TB is a fully curable and treatable disease
- Listen to the patient and assure them the services they need are available
- Refer to mental health services for further psycho-social support where required



#### References

- **1.** Das J et al, In Urban And Rural India, A Standardized Patient Study Showed Low Levels Of Provider Training And Huge Quality Gaps. *Health Affairs* 31,No. 12 (2012): 2774–2784.
- **2.** Das J et al, Use of standardised patients to assess quality of tuberculosis care: a pilot, cross-sectional study. *The Lancet*, S1473-3099(15)00077-8.
- **3.** Medicines Sans Frontieres. HIV/TB Counselling: Who is doing the job? Time of recognition of lay counsellors. URL: http://www.msf.org/sites/msf. org/files/final\_web\_counsellor\_repor t.\_one\_page.pdf
- **4.** Ministry of Health and Family Welfare, Revised National Tuberculosis Control Programme (RNTCP) Annual Status Report 2015
- **5.** Ministry of Health and Family Welfare, National Strategic Plan (2012 2017)
- 6. World Health Organisation (WHO), Standards for TB Care in India (STCI) 2014
- **7.** TB Alert. Side effects of TB drugs. URL: http://www.tbalert.org/about-tb/global-tb-challenges/side-effects/
- **8.** Hurtig AK et al, Tuberculosis control and directly observed therapy from the public health/human rights perspective. *International Journal of Tuberculosis Lung Disease*. 1999; 553-560 URL: http://www.ingentaconnect.com/ content/iuatld/ijtld/1999/
- **9.** Peltzer K et al. Prevalence of psychological distress and associated factors in tuberculosis patients in public primary care clinics in South Africa. *BMC Psychiatry*. 2012;12:89.
- **10.** Doherty AM et al . A review of the interplay between tuberculosis and mental health. *Gen Hosp Psychiatry.* 2013;35(4):398–406.
- **11.** Segerstrom SC, Miller GE. Psychological Stress and the Human Immune System: A Meta-Analytic Study of 30 Years of Inquiry. *Psychological bulletin*. 2004;130(4)
- **12.** Argiro P et al. Psychiatric Morbidity and Other Factors Affecting Treatment Adherence in Pulmonary Tuberculosis Patients, Tuberculosis Research and Treatment, vol. 2013, Article ID 489865, 37 pages, 2013.



- **13.** Sabaté E. Adherence to long-term therapies. Evidence for action. Geneva, World Health Organization, 2003. URL: http://apps.who.int/medicinedocs/pdf/s4883e/s4883e.pdf
- **14.** Central TB Division and National AIDS Control Organization. Training Modules for ICTC Counsellors on TB/HIV Coordination, New Delhi: Government of India; 2010. URL: http://www.tbcindia.nic.in/pdfs/Basic%20 Training%20Module%20for%20Counselors%20on%20HIV-TB.pdf.
- **15.** PATH. Interpersonal Communication and Counselling for TB and HIV Patients. URL: <a href="http://www.path.org/publications/files/CP">http://www.path.org/publications/files/CP</a> ukraine to hiv ipcc 3.pdf
- **16.** Providing Comprehensive, Patient-Centered Care: A Conceptual Framework Of Social Support For TB Patients. University Research Co. LLC URL: http://tbcare2.org/sites/default/files/Framework%20for%20social%20 suppor t%20of%20TB%20patients revised.pdf.
- **17.** Van den Hof S, Collins D, Leimane I, Jaramillo E, Gebhard A. Lessons Learned From Best Practices In Psycho-Socio-Economic Support For Tuberculosis Patients. WHO, USAID, MSH; 2014. Available at: https://www.msh.org/sites/msh.org/files/tbcare\_report\_best\_practices\_pyscho-socio economic support for tb patients nov 2014.pdf.
- 18. World Care Council. Patients' Charter for Tuberculosis Care 2006
- **19.** Burudpakdee, Chakkarin, Khan, Zeba, et al, Impact of Patient Programs on Adherence and Persistence in Inflammatory and Immunologic Diseases: A Meta-Analysis.Patient Preference and Adherence. Dove Medical Press, n.d. Web. 06 Oct. 2015
- **20.** International Union Against Tuberculosis and Lung Disease, Project Axshya. (Accessed October 6, 2015). URL: www.theunion.org/what-we-do/technical-assistance/tuberculosis-and-mdr-tb/project-axshya
- **21.** About Saksham. Tata Institute for Social Sciences. URL: www.tissgfat- mr7.com/hompage