



Let's Talk

TB:

A Supplement to GP
CLINICS

Chapter 9: Management of Drug-Resistant Tuberculosis: Q&A for Primary Care Physicians

Authors: Sujeet Rajan, MD—co-author
Madhukar Pai, MD, PhD—Author and Series Editor

What is MDR-TB and why does it matter?

- MDR-TB is caused by TB bacteria that are resistant to at least isoniazid (INH) and rifampicin (RIF)
 - 2 most potent TB drugs in the standard, short-course TB treatment
- According to WHO, India had an estimated 62,000 MDR-TB cases among notified pulmonary TB patients
 - This number is likely to be an underestimate as privately treated TB patients may not be notified

- Early diagnosis of MDR-TB is critical because the treatment regimen required is entirely different from standard, first-line anti-TB therapy (ATT)
- Treatment of MDR-TB is much longer (at least 2 years), more toxic, and very expensive
- Even with correct management mortality rates can approach 50%
- Specialized care is very important and treatment must be guided by drug-susceptibility testing (DST) results

What is XDR-TB and how common is it?

- **XDR-TB** is more severe resistance than MDR-TB
- **XDR-TB** occurs when TB bacteria become resistance to:
 - INH and rifampicin
 - +
 - Any fluoroquinolone (e.g. moxifloxacin or levofloxacin)
 - +
 - At least one of 3 injectable second-line drugs
 - amikacin, kanamycin, or capreomycin

- Treatment of XDR- TB should never be done in primary care settings
 - Specialist treatment is mandatory
- XDR-TB and pre-XDR TB increasingly seen in cities such as Mumbai
- Are considered rare at the national level in India

What is Totally Drug Resistant TB (TDR-TB)?

- Some hospitals (including in Mumbai) have reported a few cases where TB bacteria were found to be resistant to all of the drugs tested (first and second line)
 - Such resistant strains have also been reported in other countries
- While WHO does not recognize TDR-TB as a specific category, it is important to acknowledge the emergence of very extreme forms of DR-TB
 - NO drug therapy might work, and mortality rates are exceedingly high

When Should You Suspect Drug-Resistant TB in Your Patient?

- DR-TB is more common in patients who:
 - Do not take their TB medications regularly (i.e. in patients who are poorly adherent to ATT)
 - Do not respond to standard ATT: they may be getting adequate treatment in the right doses during the intensive treatment but show no clinical improvement
 - These patients may be persistently sputum AFB positive despite 2 months of adequately dosed drugs

- Develop TB disease again, after having taken ATT in the past (i.e. in any patient with recurrence or relapse)
- Come from areas of India (e.g. Mumbai) where drug-resistant TB is widespread
- History of exposure to a patient with pulmonary DR- TB

If you suspect DR-TB in your patient, how can you confirm it? What are the methods available for Drug Susceptibility Testing (DST)

- The WHO recently announced the post-2015 “End TB Strategy”
- A key component of this strategy is the push towards ‘universal DST’ which means that all TB patients should get a DST done at the time of their diagnosis
- At the very least all patients with history of previous TB treatment and treatment failure or recurrence must undergo DST
- In cities such as Mumbai it is important to get a DST on ALL TB patients regardless of risk factors

- DST can be done using two methods: genotypic and phenotypic
- Genotypic methods are based on molecular tests that detect mutations in TB bacteria that confer drug-resistance
 - For example, mutations in the *rpoB* gene of *M. tuberculosis* is strongly associated with rifampicin resistance
 - Genotypic tests include Xpert MTB/ RIF (GeneXpert) and Hain Genotype MTBDRplus (a commercial line probe assay)
- Phenotypic methods are based on detection of culture growth with and without TB drugs added to the culture media
 - Phenotypic methods include solid and liquid cultures
 - While solid cultures can take up to 2 months, liquid cultures (e.g. MGIT culture) can produce results within 2 weeks


- According to International Standards for TB Care (ISTC), DST should be performed at the start of therapy for all previously treated patients
- Patients who remain sputum smear-positive at completion of the intensive phase of treatment and patients in whom treatment has failed, have been lost to follow-up, or relapsed following one or more courses of treatment should always be assessed for drug resistance

- For patients in whom drug resistance is considered to be likely, an Xpert MTB/ RIF[®] test should be the initial diagnostic test, as per WHO policy
- Line-probe assay or liquid culture and DST to at least isoniazid and rifampicin should be performed promptly if rifampicin resistance is detected
- If MDR-TB is detected, DST to second line TB drugs especially fluoroquinolones and injectable drugs is required for correct management

- For DST, both WHO and ISTC recommend Xpert MTB/RIF[®] test as the initial diagnostic test
 - Can rapidly detect rifampicin resistance within 90 minutes with high accuracy
 - Allow clinicians to initiate empiric MDR therapy, pending confirmation with cultures
- In areas (e.g. Mumbai) where levels of primary drug resistance are high, it is imperative to send out the sputum (in pulmonary TB) and/or other samples from appropriate sites (in extra-pulmonary TB) for an Xpert MTB/RIF test

- Line-probe assays (e.g. Hain Genotype MTB-DR plus) or liquid culture and DST should then be performed promptly if rifampicin resistance is detected using Xpert MTB/RIF
- Once culture and DST results are obtained, MDR therapy can be customized to the patient's drug-susceptibility profile
 - Must include a combination of at least 5 drugs to which the TB bacilli are still sensitive

Algorithm for Drug-Susceptibility Testing (DST)



Is it important to always send a sample for MGIT culture?

- Yes
- Especially if you are suspecting MDR-TB in your patient, or practicing in an area of high prevalence of MDR-TB
- The liquid culture is the gold standard for the diagnosis of TB and MDR-TB
- Every other test is compared against the liquid culture

- Since culture results can take a few weeks, it is important to send out cultures early on in the treatment for the following reasons:
 - It is the gold standard for the confirmation of TB itself
 - Both pulmonary and extra-pulmonary samples can be subjected to culture
 - Culture confirms the diagnosis of TB as opposed to atypical mycobacteria or nocardiosis (smear microscopy may still be positive in the latter two)
 - A positive culture can be subjected to a line probe assay for an early diagnosis of MDR-TB
 - Cultures are the only tests that assay a variety of first and second line drugs, and confirm MDR and XDR-TB

- When you practice in a high-prevalence area of TB
 - Save a lot of time for the patient by sending out cultures early on in the treatment
- In a scenario where the patient turns out to have drug-sensitive strains on a culture you can rest assured you are on the right track
 - And the money spent on culture and first-line DSTs was worth it
- If the patient turns out drug-resistant, then you have saved an enormous amount of time of the patient on diagnosis and appropriate individualized treatment by the concerned specialist

When Should You Refer the Patient to a Specialist?

- To confirm the diagnosis of TB, if in doubt
- To obtain invasive samples for testing (e.g. bronchoalveolar lavage, or pleural biopsy)
- To get a treatment plan, if the patient has co-morbidities like diabetes, immunosuppressive conditions, etc. Remember diabetes triples the risk of developing TB, and increase the severity of TB too. Conversely TB can worsen blood glucose control as well, in patients with TB

- To decide on addition of steroids to an ATT regimen
- If adverse effects are unmanageable at primary care level
- If drug-resistant TB is confirmed (through molecular or culture tests) for a management and monitoring plan
- If lung resection surgery is indicated for any reason, be it for the diagnosis or treatment of the tuberculosis

- Patients can be referred to specialists in either the private/public sector
- The Revised National TB Control Program (RNTCP) in India is steadily increasing its capacity to offer MDR-TB therapy via Programmatic Management of Drug-resistant TB (PMDT) centers and hospitals
 - Where free MDR-TB therapy is available
 - This is a useful option for patients unable to afford private care

What are the commonest management errors that can result in acquired drug resistance?

- Addition of a single drug to a failing regimen
 - Many physicians add a quinolone to the 4 first-line drugs (HRZE) when the standard therapy does not result in improvement
 - This is wrong and believed to be one of the causes of high fluoroquinolone resistance in MDR-TB of late
- Prescribing only 2 or 3 drugs in the intensive phase
 - The intensive phase should always have 4 drugs
 - Otherwise, the bacteria mutate and develop resistance

- Prescription of second-line drugs in the first instance
 - Some physicians have the mistaken perception that second-line drugs are more potent than first-line medication
 - In fact they are less effective (and more toxic) drugs and should be reserved only for patients with DR-TB or first-line drug intolerance
- Prescribing total drugs for an inadequate duration
 - e.g. 2 to 4 months, instead of the minimum of 6 months

- Under-dosing of first-line medication is another common problem
 - Many physicians prescribe drug ‘kits’ or fixed dose combinations (FDCs) in the belief that everyone falls in a fixed weight category
 - This is obviously not the case
 - A significant number of patients with TB are significantly underweight and overweight too, and this needs appropriate adjustment of dosing

- Sometimes, patient switch between doctors, and if the new doctor changes the regimen without adequate drug-susceptibility information, then this might result in inappropriate regimens that increase risk of resistance
- Patients may feel better and decide to stop taking medications
 - Premature stoppage occurs with disastrous consequences on cure
 - And of course increased risk of drug-resistant bacilli

- Drug-related side effects (if not adequately counseled on at the outset) is another common reason for non-adherence and possible treatment default
- Stigma of the disease and sometimes cost (though not very high for first-line drugs) has been a reason for non-adherence and treatment default in the past
- Personal/social issues have also been reasons for patients interrupting their treatment at crucial stages (especially intensive phase) with resultant disastrous consequences
 - Village travel
 - Death in the family
 - Other

Is it proper to treat MDR-TB (Once Confirmed) in General Practice?

- No, this is not advisable for the following reasons:
 - MDR and XDR-TB do not have a simple, standardized treatment plan
 - Drug therapy needs to be customized to the individual patient's DST profile
 - The cost and duration of treatment are significant and need to be appropriately counseled on
 - The side effects of the various drugs are significant and common, and often difficult to manage
 - Hospitalization may be required to manage severe cases
 - A significant number of patients default on treatment, spreading the disease further
 - Hence significant time needs to be spent with the patient, something not easy in a primary care practice

- A lot of treatment is based on experience of the specialist and his ability to counsel patients and their care-givers appropriately
 - Early detection and treatment of XDR is also very critical here
- A significant number of patients may benefit with surgery in MDR-TB
 - Examples include decortications, lobectomy and pneumonectomy
 - Decisions regarding this should be taken by a respiratory physician early in the treatment course, in consultation with an experienced thoracic surgeon
 - These are decisions beyond the purview of primary care

Is it proper to monitor MDR-TB treatment in primary care?

- Absolutely yes!
- Every patient trusts his or her GP the most
- And especially if the GP is competent to manage drug-sensitive TB well, and alert to refer to a specialist when required
- These qualities in a GP increase trust by the patient

- The patient with MDR-TB will always need a primary care physician to monitor her disease
- Especially sputum cultures when appropriate, drug-related side effects, communication with the specialist on need for interval bronchoscopies (to ascertain culture status in lung MDR-TB) and decisions on major surgery
 - Your patient will value your role here

What are the second line drugs for TB and their side effects?

- **Group 1** are the first-line drugs:
 - isoniazid, rifampicin, ethambutol, pyrazinamide
- **Group 2:** Aminoglycosides: (any one drug to be used)
 - Streptomycin, Kanamycin, Amikacin, Capreomycin
 - 15 mg/kg/day
 - Monitor creatinine, vestibular and ototoxicity
 - Check for giddiness, Romberg's sign, tinnitus or reduced hearing as early signs of ototoxicity
 - Most MDR-TB patients are streptomycin resistant too, so avoid using streptomycin in an empiric MDR regimen
 - Kanamycin and Amikacin exhibit some cross-resistance
 - Capreomycin exhibits no cross-resistance
 - Patients who are unable to take regular intramuscular injections (up to 6 months at least), may need to take intravenous amikacin through a PICC line

- **Group 3**

- *Fluoroquinolones (ofloxacin, levofloxacin and moxifloxacin)* – any one to be used

- Usually cross-resistant to each other, except for moxifloxacin which can still exhibit activity despite resistance to other fluoroquinolones
 - Giddiness, headache and GI upsets are the commonest side effects
 - Teratogenic effects
 - Concern in small children with growing cartilage

- **Group 4** – Less potent oral agents
 - *Thioamides (ethionamide and prothionamide)*
 - 15 – 20 mg/kg/day (max 1 gm/day)
 - Epigastric discomfort – hence start with 250 mg twice daily initially and then increase the dose if tolerated
 - Hypothyroidism
 - Hypoglycaemia
 - Teratogenic effects
 - Cross-resistance between drugs

– *Cyloserine and Terizidone*

- 10- 20 mg/kg/day
- Altered behavior, mood swings, suicidal tendencies
- Giddiness, seizures
- Avoid in alcoholics, epileptics and mental illness
- Slurred speech

– *Para-aminosalicylic acid (PAS)*

- 150mg/kg/day in 2 divided doses
- GI upsets
- Avoid the sodium salt in patients needing salt restriction
- Hypothyroidism

- **Group 5** – Drugs for whom anti-tuberculosis action has not been documented in clinical trials
 - Thioacetazone
 - Clofazimine
 - Linezolid
 - Amoxicillin-clavulanic acid
 - Clarithromycin
 - Imipenam – cilastatin
 - High dose isoniazid

New TB drugs for MDR-TB that are not approved for clinical use in India

- 2 new drugs are now available for TB:
 - Bedaquiline and Delamanid
 - Clinical trials on these drugs are ongoing and approval by the Indian regulatory agency is expected in future
 - Currently, these drugs should NOT be used by primary care providers
 - They might be available for specialist use on compassionate grounds

Bedaquiline

- SIRTURO® by Janssen Therapeutics TM
- Bedaquiline (<https://www.sirturo.com/>) is a diarylquinoline anti mycobacterial drug indicated as part of combination therapy in adults (≥ 18 years) with pulmonary multi-drug resistant tuberculosis (MDR-TB)
- SIRTURO was the first new TB drug with a novel mechanism of action to be made available for more than 40 years and was granted accelerated approval by the United States FDA
- Still in Phase III trials
- SIRTURO is restricted for use only when an effective treatment regimen cannot otherwise be provided

Delamanid

- Delamanid (also known by its trade name, Deltyba® by Otsuka Pharmaceutical Co. Ltd) is a drug in the nitroimidazole class
- Delamanid is a new potential option for people with MDR-TB who lack effective, tolerable treatments
- Delamanid was granted conditional approval by the European Medicine Agency in April 2014
- Information about this new drug is limited
 - Has only been through Phase IIb trial and studies for safety and efficacy
- Interim policy guidance on these new drugs is available from WHO

What are the things to remember when checking the prescription of a specialist for MDR-TB?

- As a GP, your **role is paramount**
- A good primary care physician can be alert to wrong prescriptions and remind a specialist at times, that the direction or plan of treatment is wrong
- Remember you see the patient far more frequently than the specialist
- Check that at least 4 new drugs have been commenced
 - Preferably one from Group 2 and 3
 - 2 more from **Group 3, 4** or **both** (depending on whether the patient has MDR, XDR or TDR TB)

- Check that none of the drugs are contraindicated for your patient
 - Hearing loss pre-existing
 - Mental Illness
 - Visual Disturbances
 - Chronic Kidney Disease
 - Pre-existing seizure disorder
 - Pre-existing severe gastritis

- Do a skin sensitivity test before administering the aminoglycoside the first time
- A drug that was used within a previous failing regimen should never be used again if possible
 - Should not be counted in the total of 4 drugs for re-treatment
- Monitor CBC and platelets in patient on linezolid
- Monitor for visual disturbances in patient on linezolid

Questions you need to ask the specialist periodically during the treatment of MDR-TB

- Is the patient non-infectious?
- When do you next need a sputum/bronchoscopic sample for AFB culture?
- When should the next round of chest x-rays be done?
- Is a CT scan chest indicated, and if so, for what purpose?
- If there is a persistent cavity despite clinical improvement, how can we ensure the patient is still not harboring active infection? (usually a bronchoscopic aspirate is indicated here)
- What is the likely total cost of treatment, excluding any surgery?
- Is the patient a candidate for surgery, and if so what is the likely cost?
- How long do you expect the entire duration of treatment to last?

Importance of asking these questions

- It is very important to ask these questions
- The specialist's message needs to be constantly reinforced by the primary care physician
 - Ensures better adherence at all stages of treatment

Treating and Supervising MDR-TB Management

- Treating and supervising MDR-TB management in clinical practice is a big responsibility and needs commitment and time
- If you are unable to manage the same, refer to an appropriate specialist
- **NEVER feel you will lose your patient that way!**
 - Your role to monitor for side effects is paramount; so too your role in ensuring adherence
 - Patients are ever-grateful when they are referred to the right specialists
 - Their confidence in you (as their primary care physician) will only increase

References

1. World Health Organization. Global Tuberculosis Report 2014. Geneva: WHO, 2014.
2. Udwadia ZF. MDR, XDR, TDR tuberculosis: ominous progression. *Thorax*. 2012; 67(4): 286-8.
3. Udwadia ZF, Amale RA, Ajbani KK, Rodrigues C. Totally drug-resistant tuberculosis in India. *Clin Infect Dis*. 2012; 54(4): 579-81.
4. Dalal A, Pawaskar A, Das M, et al. Resistance Patterns among Multi- drug-Resistant Tuberculosis Patients in Greater Metropolitan Mumbai: Trends over Time. *PLoS One*. 2015; 10(1): e0116798.
5. World Health Organization. The End TB Strategy. Global strategy and targets for tuberculosis prevention, care and control after 2015. URL: http://www.who.int/tb/post2015_TBstrategy.pdf?ua=12014. (accessed 27 Oct 2014).
6. TB CARE I. International Standards for Tuberculosis Care, 3rd Edition. URL: www.istcweb.org 2014. (accessed March 2014).
7. World Health Organization. Policy update: automated real-time nucleic acid amplification technology for rapid and simultaneous detection of tuberculosis and rifampicin resistance: Xpert MTB/RIF system for the diagnosis of pulmonary and extrapulmonary TB in adults and children 2013. <http://www.stoptb.org/wg/gli/assets/documents/WHO%20Policy%20Statement%20on%20Xpert%20MTB-RIF%202013%20pre%20publication%2022102013.pdf> (accessed).
8. WHO. Guidelines for the programmatic management of drug-resistant tuberculosis. URL: http://www.who.int/tb/challenges/mdr/programmatic_guidelines_for_mdrtb/en/ (accessed 8 August 2015).